

# INDEX

<b>Sr. No.</b>	<b>Topics</b>	<b>Page No.</b>
1	Declaration	ii
2	Certificate	iii
3	Certificate	iv
4	Copyright	v
5	Acknowledgement	vi
6	Preface	ix
7	Index	xi
8	List of Tables	xiv
9	List of Figures	xix
10	List of Abbreviations	xx
<b>1</b>	<b>INTRODUCTION</b>	<b>1 – 35</b>
<b>2</b>	<b>LITERATURE REVIEW</b>	<b>36 – 49</b>
<b>3</b>	<b>RESEARCH GAP</b>	<b>50 – 51</b>
<b>4</b>	<b>AIM,OBJECTIVES, PLAN OF WORK</b>	<b>52 – 53</b>
<b>5</b>	<b>MATERIAL AND METHOD</b>	<b>54 – 106</b>
<b>6</b>	<b>RESULT AND DISCUSSION</b>	<b>107 – 182</b>
<b>7</b>	<b>SUMMARY AND CONCLUSION</b>	<b>183 – 184</b>
<b>8</b>	<b>REFERENCES</b>	<b>185 - 200</b>
	<b>APPENDIX:</b>	
	Research and Review paper publication	-
	Conference Certificates	-

## LIST OF TABLES

Table No.	Title of the Table	Page No.
1.1	List of chemical drugs used as Antidiabetic	28
5.1	Chemical constituents present in <i>Sesbania Grandiflora</i> leaves	58
5.2	Different types of <i>Beta Vulgaris</i> L	65
5.3	Description of <i>Beta vulgaris</i> L	67
5.4	Chemical constituent present in <i>Beta Vulgaris</i> root	67
5.5	Description of Magnesium Stearate	70
5.6	Description of Talc	71
5.7	Description of Lactose Anhydrous	72
5.8	Description of Microcrystalline Cellulose	73
5.9	List of Materials used	74
5.10	List of Instruments /Equipments used	76
5.11	Chemical test for detection of Alkaloids.	85
5.12	Chemical test for detection of Carbohydrates.	86
5.13	Chemical test for detection of Glycosides.	86
5.14	Chemical test for detection of Protein.	87
5.15	Chemical test for detection of Phytosterols.	87
5.16	Chemical test for detection of Fixed oils and Fats	88
5.17	Chromatographic conditions	93
5.18	Drug combination ratio	94
5.19	Composition of Polyherbal formulation.	101
5.20	Flow properties and corresponding Angle of Repose	102

5.21	Grading of powders for their flow properties	103
5.22	Compressibility Index and Hausner's Ratio for powder flow	104
5.23	The Limits of the weight variation	105
6.1	Description of <i>S. Grandiflora</i> L	110
6.2	Description of <i>Beta vulgaris</i> Root.	111
6.3	Ultra-Violet analysis of leaves <i>Sesbania Grandiflora</i>	114
6.4	Ultra-Violet analysis of root of <i>Beta Vulgaris</i>	114
6.5	Angle of Repose of <i>Sesbania Grandiflora</i> leaves powder and <i>Beta Vulgaris</i> Root powder	116
6.6	Physicochemical Parameters of leaves <i>Sesbania Grandiflora</i> and root of <i>Beta Vulgaris</i> L	118
6.7	Extraction of leaves of <i>Sesbania Grandiflora</i> and root of <i>Beta vulgaris</i> L	119
6.8	Qualitative Phytochemical Tests of leaves <i>Sesbania Grandiflora</i>	122
6.9	Qualitative Phytochemical Tests of root of <i>Beta Vulgaris</i>	124
6.10	pH of Extracts	126
6.11	Absorbance of Gallic acid at 760 nm	126
6.12	Absorbance of Quercetin at 510 nm	128
6.13	Total phenolic and flavonoid content of leaves extracts of <i>Sesbania Grandiflora</i>	129
6.14	Total phenolic and flavonoid content of leaves extracts of <i>Beta Vulgaris</i>	129
6.15	Functional group in methanolic extract of <i>Sesbania Grandiflora</i> leaves	130
6.16	Functional group in methanolic extract of <i>Beta Vulgaris</i> Root	131
6.17	HPTLC fingerprinting of Methanolic extract of root of <i>Beta Vulgaris</i> at 254 nm and 366 nm	133

6.18	Rf Value of Methanolic extract of root of <i>Beta Vulgaris</i>	133
6.19	Peak table of Methanolic extract of root of <i>Beta Vulgaris</i> with Rf Values at 254 nm and 366 nm	133
6.20	Peak table of Betalain of root of <i>Beta Vulgaris</i> with Rf Values at 254 nm and 366 nm	135
6.21	HPTLC fingerprinting of Methanolic extract of leaves of <i>Sesbania Grandiflora</i> of at 254 nm and 366 nm	136
6.22	Rf Value of Methanolic extract of leaves of <i>Sesbania Grandiflora</i>	136
6.23	Peak table of Methanolic extract of leaves of <i>Sesbania Grandiflora</i> with Rf Values at 254 nm and 366 nm	137
6.24	Peak table of Kaepferol and Quercetin of leaves of <i>Sesbania Grandiflora</i> with Rf Values at 254 nm and 366 nm	138
6.25	Observation of changes in Clinical Signs in PHF (2000 mg/kg) Administered Acute Toxicity Group	141
6.26	Histopathology of Pancreas in groups	166
6.27	Histopathology of Liver in groups	170
6.28	Composition of Polyherbal Formulation	174
6.29	Organoleptic characters of Polyherbal Formulation	176
6.30	Functional group in Polyherbal Formulation	177
6.31	HPTLC fingerprinting of Methanolic extract combination in equal ratio at 254nm and 366nm	178
6.32	R <sub>f</sub> Value of Methanolic extract of root of PHF2	178
6.33	Preformulation studies	179
6.34	Results of Post compression studies of Polyherbal Formulation	180
6.35	Results of Stability study of F3 Polyherbal Tablet	185

## LIST OF FIGURES

Figure No.	Title of the Figure	Page No.
1.1	Herbal Medicine used in diseases	03
1.2	Standardization of Herbal Drug	11
1.3	Major Endocrine glands, Tissues and their hormones	17
1.4	Type 1 Diabetes	18
1.5	Type 2 Diabetes	19
1.6	Islets of Langerhans	21
1.7	Secretion of Insulin in $\beta$ cells	22
5.1	<i>Sesbania Grandiflora</i> L plants	54
5.2	<i>Beta Vulgaris</i> L.Root	64
5.3	Determination of Loss on drying (Moisture Content)	80
5.4	Alcohol Soluble Extractive Value	80
5.5	Water Soluble Extractive Value	81
5.6	Furnace of Ash value	82
5.7	Extraction of <i>Sesbania Grandiflora</i> and <i>Beta vulgaris</i> powder	84
5.8	Lyophilizer	85
5.9	Preliminary Phytochemical screening of Plant Extract	89
5.10	pH meter	89
5.11	FTIR spectroscopy	91
5.12	OECD guidelines for Acute toxicity studies	96
5.13	<i>Invivo</i> studies	99
5.14	Tablet compression machine	100
6.1	Authentication letters of Drugs	108
6.2	<i>Sesbania Grandiflora</i> leaves powder and <i>Beta Vulgaris</i> root	109

	powder	
6.3	<i>Sesbania Grandiflora</i> Leaves	110
6.4	<i>Beta Vulgaris</i> L Root	110
6.5	Microscopic Evaluation of <i>Sesbania Grandiflora</i> Linn	112
6.6	Microscopic Evaluation of <i>Beta Vulgaris</i> Root	113
6.7	Angle of Repose of <i>Sesbania Grandiflora</i> leaves powder and <i>Beta Vulgaris</i> Root powder	115
6.8	Alcohol soluble Extractive	117
6.9	Water Soluble Extractive	117
6.10	Rotary Evaporator	119
6.11	Aqueous Extract leaves of <i>Sesbania Grandiflora</i> and root of <i>Beta vulgaris</i> L	119
6.12	Methanolic Extract of leaves of <i>Sesbania Grandiflora</i> and root of <i>Beta vulgaris</i> L	120
6.13	Acetone Extract of leaves of <i>Sesbania Grandiflora</i> and root of <i>Beta vulgaris</i> L	120
6.14	Ethanollic extract of leaves of <i>Sesbania Grandiflora</i> and root of <i>Beta vulgaris</i> L	120
6.15	Standard curve of Gallic acid at 760 nm	128
6.16	Standard curve of Quercetin at 510 nm	128
6.17	I.R Spectra of methanolic extract of <i>Sesbania Grandiflora</i> leaves	130
6.18	I.R Spectra of methanolic extract of <i>Beta Vulgaris</i> root	131
6.19	HPTLC fingerprinting of Gallic acid and Quercetin at 254 nm and 366 nm	132
6.20	HPTLC fingerprinting of Methanolic extract of root of <i>Beta Vulgaris</i> at 254 nm and 366 nm	134
6.21	HPTLC Densitogram of Methanolic root Extract Of <i>Beta Vulgaris</i>	135

6.22	HPTLC Densitogram of Methanolic extract of leaves of <i>Sesbania Grandiflora</i> at 254 nm and 366 nm	137
6.23	Densitogram of Methanolic Leaves Extract Of <i>Sesbania Grandiflora</i>	138
6.24	Animal Ethical Letter for <i>In vivo</i> Studies for Antidiabetic Activity.	140
6.25	Effect of PHF 1, PHF 2, PHF 3 extract on blood glucose level (mg/dl)	141
6.26	Metabolic cages for Urine collection	141
6.27	Microscopical examination of Urine collected after treatment of PHF1, PHF2, PHF3	142
6.28	Effects of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on the red blood cell (RBC) count (Million/Cu mm) in STZ-induced diabetes in rats.	143
6.29	Effects of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on haemoglobin levels (g/dl) in rats with streptozotocin-induced diabetes.	144
6.30	Effect of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on packed cell volume (PCV) in rats with streptozotocin -induced diabetes.	145
6.31	Effects of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on mean corpuscular volume (MCV, fl) in rats with streptozotocin-induced diabetes.	145
6.32	Effects of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on mean corpuscular hemoglobin (MCH) in rats with streptozotocin-induced diabetes.	146
6.33	Effect of PHF2 on mean corpuscular hemoglobin concentration (MCHC) in rats with diabetes induced by	147

	streptozotocin.	
6.34	Effect of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on white blood cell (WBC) count in rats with streptozotocin-induced diabetes.	147
6.35	Effect of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on polymorphonuclear leukocytes (polymorphs) in rats with streptozotocin-induced diabetes.	148
6.36	Effect of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on lymphocyte levels in streptozotocin-induced diabetic rats.	149
6.37	Effect of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on eosinophil levels in rats with streptozotocin-induced diabetes.	149
6.38	Effects of PHF2 administered at doses of F200 and F400 on monocyte levels in rats with streptozotocin-induced diabetes are illustrated.	150
6.39	Effects of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on basophil counts in streptozotocin-induced diabetic rats.	151
6.40	Effect of PHF2 at doses of F200 and F400 on platelet count (Lakhs/Cumm) in rats with streptozotocin-induced diabetes.	151
6.41	Effect of PHF2 administration on blood glucose levels in rats with streptozotocin-induced diabetes.	152
6.42	Effect of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on creatinine levels (mg/dl) in rats with streptozotocin-induced diabetes.	153



6.43	Effect of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on serum protein levels (mg/dl) in rats with streptozotocin-induced diabetes.	154
6.44	Effects of PHF2 at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on alanine transaminase (ALT) levels (IU/L) in rats with streptozotocin-induced diabetes.	155
6.45	Effects of PHF2 at doses of F200 and F400 on aspartate transaminase levels (IU/L) in rats with streptozotocin-induced diabetes	156
6.46	Effect of PHF2 at doses F200 and F400 on blood urea nitrogen (BUN) levels (mg/dl) in rats with streptozotocin-induced diabetes.	157
6.47	Effects of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on total cholesterol levels (mg/dl) in rats with streptozotocin-induced diabetes.	158
6.48	Effect of PHF2 administered at doses of F200 and F400 on triglyceride levels (mg/dl) in rats with streptozotocin-induced diabetes is depicted.	159
6.49	Effect of PHF2 at doses of F200 and F400 on high-density lipoprotein (HDL) levels (mg/dl) in rats with streptozotocin-induced diabetes.	160
6.50	Effect of PHF2 given at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on low-density lipoprotein (LDL) levels (mg/dl) in rats with streptozotocin-induced diabetes.	161
6.51	Effects of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on very low-density lipoprotein (VLDL) levels (mg/dl) in rats with streptozotocin (STZ)-induced diabetes.	162
6.52	Effects of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on Glucose absorption levels mM in rats with streptozotocin (STZ)-induced	163

	diabetes.	
6.53	Effects of PHF2 administered at doses of 200 mg/kg (F200) and 400 mg/kg (F400) on HbA1c level in rats with streptozotocin (STZ)-induced diabetes.	164
6.53	Histopathology of Pancreases in group	166
6.54	Histopathology of Liver in group	171
6.55	F1 powder	174
6.56	F2 powder	174
6.57	F3 powder	175
6.58	F3 powder	175
6.59	F1 to F4 Tablets	175
6.60	FTIR of Polyherbal formulation	176
6.61	HPTLC finger printing of Methanolic extract of root of PHF2 at 254 nm and 366 nm	178
6.62	Angle of repose PHF powder	179
6.63	Bulk and Tapped Density of PHF	180
6.64	Average Weight of Tablet	181
6.65	Hardness of Tablet.	181
6.66	Thickness of Tablet.	181
6.67	Friability test of Tablet.	181
6.68	Disintegration test of Tablet.	181

## LIST OF ABBREVIATIONS

<b>SG</b>	: <i>Sesbania Grandiflora</i>
<b>BV</b>	: <i>Beta Vulgaris</i>
<b>WHO</b>	: World Health Organization
<b>DM</b>	: Diabetes Mellitus
<b>MODY</b>	: Maturity-onset diabetes of the young
<b>(FPG)</b>	: Fasting plasma glucose
<b>PHF</b>	: Polyherbal Formulation
<b>T1DM</b>	: TYPE 1 Diabetes Mellitus
<b>T2DM</b>	: TYPE 2 Diabetes Mellitus
<b>BGL</b>	: blood glucose levels
<b>HFD</b>	: high-fat diet
<b>HNF</b>	: Hepatocyte nuclear factor
<b>SGOT</b>	: Serum glutamic pyruvic transaminase
<b>SGPT</b>	: Serum glutamic oxaloacetic transaminase
<b>HPTLC</b>	: High-Performance Thin Layer Chromatographic
<b>BUN</b>	: Blood urea nitrogen
<b>STZ</b>	: Streptozotocin
<b>ANOVA</b>	: Analysis Of Variance
<b>Rf</b>	: Reference Factor
<b>M.P</b>	: melting point
<b>NIDDM</b>	: Non-Insulin Dependent Diabetes Mellitus
<b>GD</b>	: Gestational Diabetes
<b>UV-Vis</b>	: Ultra violet Visible
<b>FT-IR</b>	: Fourier Transform Infra-Red
<b>TLC</b>	: Thin Layer Chromatography
<b>HPLC</b>	: High Performance Liquid Chromatography
<b>ppm</b>	: parts per million